arXiv:1405.4488v1 [math.LO] 18 May 2014

A global approach to AECs

Hugo L. Mariano *† Pedro H. Zambrano ‡ Andrés Villaveces

December 2013

Abstract

In this work we present some general categorial ideas on Abstract Elementary Classes (AECs) , inspired by the totality of AECs of the form $(Mod(T), \preceq)$, for a first-order theory T: (i) we define a natural notion of (funtorial) morphism between AECs; (ii) explore the following constructions of AECs: "generalized" theories, pullbacks of AECs, (Galois) types as AECs; (iii) apply categorial and topological ideas to encode model-theoretic notions on spaces of types ; (iv) present the "local" axiom for AECs here called "local Robinson's property" and an application (Robinson's diagram method); (v) introduce the category *AEC* of Grothendieck's gluings of all AECs (with change of basis); (vi) introduce the "global" axioms of "tranversal Robinson's property" (TRP) and "global Robinson's property" (GRP) and prove that TRP is equivalent to GRP and GRP entails a natural version of Craig interpolation property.

1 Introduction

The notion of Abstract Elementary Class (AEC) was introduced by Shelah in 1987 ([**She**]) and is a an abstract model-theoretic setting (i.e., there is no syntax and proof theory associated to it) that generalizes the natural occurrences:
(i) $T \subseteq L_{\omega,\omega} - sent \rightsquigarrow (Mod(T), \preceq)$; (ii) $\theta \in L_{\omega_1,\omega} - sent \rightsquigarrow (Mod(\theta), \preceq)$

Definition 1.1. *An AEC is a pair $(\mathcal{K}, \preceq_K)$ where $\mathcal{K} \subseteq L-str$, for some (finitary) language L and $\preceq_K$ is a binary relation on $\mathcal{K}$ that satisfies:*

(i) order*:*
- $\preceq_K$ *is a partial order relation on* $\mathcal{K}$*;*
- $M \preceq_K N \Rightarrow M \subseteq_{L-str} N$.

(ii) isomorphism*:*
- $M \in \mathcal{K}$ *and* $M' \cong_{L-str} M$*, then* $M' \in \mathcal{K}$*;*
- $M \preceq_K N$*,* $f \in L - iso(N, N')$ *and* $f \restriction \in L - iso(M, M') \Rightarrow M' \preceq_K N'$

(iii) coherence*:*
$M, N, P \in \mathcal{K}$*,* $M \subseteq_{L-str} N \subseteq_{L-str} P$ *and* $M \preceq_K P$*,* $N \preceq_K P \Rightarrow M \preceq_K N$

(iv) reunion*: Let* $\{M_\alpha : \alpha \in \theta\} \subseteq \mathcal{K}$*,* $M_\alpha \preceq_K M_\beta$ *if* $\alpha < \beta < \theta$*, then:*
- $M := \bigcup_{\alpha\in\theta} M_\alpha \in \mathcal{K}$ *and* $M_\alpha \preceq_K M$*,* $\forall \alpha \in \theta$*;*
- *if* $N \in \mathcal{K}$ *and* $M_\alpha \preceq_K N$*,* $\forall \alpha \in \theta$*, then* $M \preceq_K N$.

(v) Löwenhiem-Skolem*: There is an infinite cardinal* κ *such that for each* $N \in \mathcal{K}$ *and* $X \subseteq |N|$*, there is* $M \in \mathcal{K}$*,* $M \preceq_K N$*,* $X \subseteq |M|$ *and* $\sharp |M| \leqslant \sharp X + \kappa$*.* $LS(\mathcal{K}) := min\{\kappa$*:* κ *satisfies the property*$\}$ □

1.2. Let $(\mathcal{K}, \preceq_K)$ be an AEC with $\mathcal{K} \subseteq L - str$, then there are two natural notions:
- $\mathcal{K}$-substructure: $C \preceq_K D \Rightarrow C \overset{\preceq_K}{\hookrightarrow} D$
- $\mathcal{K}$-embedding: $(A \overset{f}{\rightarrowtail} B) = (A \overset{\cong_L}{\to} A' \overset{\preceq_K}{\hookrightarrow} B)$ □

1.3. *An AEC as a category:* Let $cat(\mathcal{K})$ be the category with $\mathcal{K}$ as the class of objects and $\mathcal{K}$-embeddings as arrows.

- This is well-defined:

identities: because $\preceq_K$ is reflexive
composition: by the closure under isos

*Research supported by FAPESP, under the Thematic Project LOGCONS: Logical consequence, reasoning and computation (number 2010/51038-0).

†Instituto de Matemática e Estatística, University of São Paulo, Brazil, email: hugomar@ime.usp.br

‡Departamento de Matemticas, Faculdad de Ciencias, National University of Colombia, Colombia, emails: phzambranor@unal.edu.co, avillavecesn@unal.edu.co

$$A \overset{\cong}{\to} A' \overset{\prec}{\hookrightarrow} B \overset{\cong}{\to} B' \overset{\prec}{\hookrightarrow} C$$
$$A' \overset{\cong}{\to} A'' \overset{\prec}{\hookrightarrow} B'$$

- $cat(\mathcal{K})$ has all (upward) directed colimits.
- $cat(\mathcal{K})$ is an accessible category: [**AR**], [**Lie1**] [**Lie2**], [**BR**].
- We will denote $cat(\mathcal{K})_{strict}$ the category with $\mathcal{K}$ as the class of objects and $\prec_K$-inclusions as arrows. □

1.4. *Shelah's Presentation Theorem* ([**She**], [**Bal2**]): Is the first occurence of "change of languages" in AECs-theory:
$\mathcal{K} \subseteq L - str \quad \Rightarrow \quad \exists L' \supseteq L \; \exists T' \subseteq L' - sent \; \exists$ a set Γ' of (finitary) L'-types such that, if $EC(T', \Gamma') = \{M' \in L' - str : M' \vDash T'$ and M' omits $\Gamma'\}$, then:
– $\mathcal{K} = \{M' \restriction_L \in L - str : M' \in EC(T', \Gamma')\}$ (i.e. $\mathcal{K} = PC_L(T', \Gamma')$);
– $M' \prec_{L'_{\omega\omega}} N' \in EC(T', \Gamma') \; \Rightarrow M' \restriction_L \prec_K N' \restriction_L \in \mathcal{K}$ □

1.5. *Main Question: Shelah's categoricity conjecture* ([**Bal1**]): There is a cardinal $\mu(\kappa)$ such that for all AEC $\mathcal{K}$ with $LS(\mathcal{K}) \leqslant \kappa$, if $\mathcal{K}$ is categorical in some cardinal greater than $\mu(\kappa)$ then K is categorical in all $\lambda \geqslant \mu(\kappa)$. □

1.6. *AECs and infinitary languages:* $LS(\mathcal{K}) \leqslant \kappa \quad \Rightarrow \quad \mathcal{K}$ is closed under $\equiv$ in L_{∞,κ^+} ([**Kue**]) □

1.7. *(Generalized) types in AECs:* The notion of type in AECs-theory ([**Lie1**], [**Lie2**]) is given by another application of "change of languages". We consider here an expansion of this notion, called "generalized type" or simply g-type (see, for instance, section I.4 in [**Van**] for motivations).

g-types: $M \in \mathcal{K}$, I set, a (generalized) I-type over M:
is an equivalence class $(N, f, \bar{a})$ where $N \in \mathcal{K}$, $f : M \rightarrowtail N \in cat(\mathcal{K})$, $\bar{a} : I \to N$;
$(N, f, \bar{a}) \sim (N', f', \bar{a}')$ iff $\exists n \in \mathbb{N}$, $P_0, \ldots, P_n \in \mathcal{K}$ and $\exists$ zigzag
$(N, f, \bar{a}) = (P_0, f_0, \bar{a}_0) \overset{h_1}{\to} (P_1, f_1, \bar{a}_1) \overset{h_2}{\leftarrow} (P_2, f_2, \bar{a}_2) \overset{h_3}{\to} \ldots \overset{h_{n-1}}{\leftarrow} (P_{n-1}, f_{n-1}, \bar{a}_{n-1}) \overset{h_n}{\to} (P_n, f_n, \bar{a}_n) = (N', f', \bar{a}')$
- LS-axiom $\Rightarrow \exists$ a set of representatives for I-types over M;
- if $cat(\mathcal{K})$ enjoys (AP), then $(N, f, \bar{a}) \sim (N', f', \bar{a}')$ iff $\exists P \in \mathcal{K}$ and morfisms h_1, h_2 such that:
$(N, f, \bar{a}) \overset{h_1}{\to} (P, g, \bar{b}) \overset{h_2}{\leftarrow} (N', f', \bar{a}')$
- if $cat(\mathcal{K})$ enjoys (AP), (JEP), has arbitrary large models, then:
– $\exists$ monster models;
– I-types over $M \leftrightsquigarrow$ orbits of I-sequences over a $\mathcal{K}$-monster model over $M \cup I$ (Galois g-types).
notation: $g - T(M, I)$
- *topology on g-types*: this is analogous to the topology on usual types considered in [**Lie1**]
$\lambda \geqslant LS(\mathcal{K})$, $M \in \mathcal{K}$:
$\mathcal{K}^\lambda(M) := \{N \in \mathcal{K} : N \prec_K M, \sharp N \leqslant \lambda\}$,
basis of $X^\lambda(M, I)$, the λ-topology of $g - T(M, I)$: $\{U_{p,N} : p \in g - T(N, I), N \in \mathcal{K}^\lambda(M)\}$, where:
$U_{p,N} := \{q \in g - T(M, I) : q \restriction = p\}$
(remark: if $\sharp M \leqslant \lambda$, then $X^\lambda(M, I)$ is discrete) □

2 The category of all AECs

2.1. *Motivation*: change of languages in model theory of FOL (**Institution theory**)
$\alpha : L \to L'$ (morphism of languages) $\quad \rightsquigarrow \quad (\hat{\alpha}, \alpha^\star)$, where:
- $\hat{\alpha} : Form(L) \to Form(L')$, a function defined by recursion on complexity
- $\alpha^\star : L' - str \to L - str$, a functor
$|M'^\alpha| = |M'|$, $R^{M'^\alpha} = \alpha(R)^{M'}$, $f^{M'^\alpha} = \alpha(f)^{M'}$, $c^{M'^\alpha} = \alpha(c)^{M'}$
- $M'^\alpha \vDash_L \theta[\bar{a}] \Leftrightarrow M' \vDash_{L'} \hat{\alpha}(\theta)[\bar{a}]$
- $f' : M' \to N'$ is L'-homomorfism (respect. L'-embbeding, L'-elementary embedding) $\quad \Rightarrow$
$f' : M'^\alpha \to N'^\alpha$ is L'-homomorfism (respect. L'-embbeding, L'-elementary embedding)
- $T \subseteq L - sent$, $T' \subseteq L' - sent$ are closed theories such that $\hat{\alpha}^{-1}[T'] \supseteq T$, then $\alpha^\star \restriction : (Mod(T'), \prec_{L'}) \to (Mod(T), \prec_L)$ □

2.2. *The category of all AECs* (For foundational issues $\rightsquigarrow$ Grothendieck's universes)
objects: the "class of all" AECs
arrows: $\mathcal{K} \subseteq L - str$, $\mathcal{K}' \subseteq L' - str$ (remark that the language is determined from the (non-empty) class, set $\emptyset \subseteq \emptyset - str$)
$\alpha : L \to L'$ such that the restriction $\alpha^\star \restriction : (\mathcal{K}', \prec_{K'}) \to (\mathcal{K}, \prec_K)$ is well defined:



composition: $(\beta \circ \alpha)^\star = \alpha^\star \circ \beta^\star$, ...

identities: ok

Example: $inclusion : (K, \preceq_K) \hookrightarrow (L - str, \subseteq_L)$, $inclusion = (id_L)^\star\restriction$.
Remark: $\alpha^\star$ is always a faitful functor. If α is a "surjective" morphism of languages, then $\alpha^\star$ is a full functor. □

2.1 Constructions of AECs

In this subsection we provide new examples of AECs, obtained from construction, for instance: elementary class (Mod(T)) in AECs; pullbacks of AECs; g-types in AECs, etc.

Theories Mod(T) in AEC

2.3. (**FOL**) The L-satisfaction relation $\models_L$:
• establish a Galois connection (Mod, Th) between sets of (first-order) L-sentences and classes of L-structures;
• $(Mod\restriction, Th\restriction)$ establish (order reversing) bijection between closed L-theories and elementary classes in language L. □

Definition 2.4. *In an AEC $\mathcal{K}\subseteq L - str$, define:*
(i) the "generalized elementary relation" in $\mathcal{K}$: $\equiv_K$ is "connection relation in $cat(\mathcal{K})$", i.e., is the equivalence relation generated by the arrows in $cat(\mathcal{K})$:
$M \equiv_K N$ *iff* $\exists n \in \mathbb{N}$, $P_0, \dots P_n \in \mathcal{K}$ *and* $\exists$ *zigzag* $M = P_0 \xrightarrow{f_1} P_1 \xleftarrow{f_2} P_2 \xrightarrow{f_3} \dots \xleftarrow{f_{n-1}} P_{n-1} \xrightarrow{f_n} P_n = N$
(ii) a subclass $\mathcal{M}\subseteq\mathcal{K}$ is a "generalized elementary subclass" or a "generalized (closed) theory", when $\mathcal{M}$ is closed under $\equiv_K$;
(iii) there is only a set (of representatives) of closed theories (by LS axiom);
(iv) a "maximal consistent g-theory" is an equivalence class module $\equiv_K$ (it is a closed theory) $(Max(\mathcal{K}) = \mathcal{K}/\equiv_K)$;
(v) a "complete g-theory" is a non-empty subset of a maximal consistent g-theory. □

Fact 2.5. *$\mathcal{M}\subseteq\mathcal{K}$ is g-elementary subclass $\Rightarrow$ $(\mathcal{M}, (\preceq_K)\restriction_M)$ is an AEC, a subAEC of $\mathcal{K}$, and $LS(\mathcal{M}) \leqslant LS(\mathcal{K})$.* □

Fact 2.6. *Let $\alpha : L \to L'$, $\alpha^\star\restriction : \mathcal{K}' \to \mathcal{K}$:*
(i) $M' \equiv_{K'} N' \Rightarrow M'^{\alpha} \equiv_K N'^{\alpha}$
(ii) Induced Theory: Let $\alpha^\star\restriction : \mathcal{K}' \to \mathcal{K}$, $Mod(T')\subseteq\mathcal{K}'$, then exists $Mod(T')^\alpha :=$ the least closed theory of $\mathcal{K}$ that contains $\{M'^{\alpha} : M' \in Mod(T')\}$ (iii) a commutative diagram:



$Mod(T')\subseteq\mathcal{K}'$ *maximal consistent theory* $\Rightarrow$ $Mod(T')^\alpha\subseteq\mathcal{K}$ *is a complete theory*

□

Limits of AECs

We will show that the category of all AECs has all finite limits because it has terminal object and pulbacks.

2.7. *The terminal AEC:* is obtained taking:
• $L = \emptyset$, $\mathcal{K} = L - str = Set$
• $M \preceq_K N \Leftrightarrow M\subseteq N$ □

2.8. *Pullbacks of AECs:*
• L, L_0, L_1
• $\alpha_i : L \to L_i$, $\alpha_i^\star\restriction : (\mathcal{K}, \preceq_K) \to (\mathcal{K}_i, \preceq_{K_i})$
• $L' = (L_0 \sqcup L_1)/\sim$

$\pi_i : L_i \to L'$ $(= L_i \rightarrowtail L_0 \sqcup L_1 \twoheadrightarrow L')$
$\delta = \pi_0 \circ \alpha_0 = \pi_1 \circ \alpha_1 : L \to L'$
$\mathcal{K}' \subseteq L' - str$
– $M' \in \mathcal{K}' \;\Leftrightarrow\; M'^{\pi_0} \in \mathcal{K}_0, M'^{\pi_1} \in \mathcal{K}_1, ((M'^{\pi_0})^{\alpha_0} = (M'^{\pi_1})^{\alpha_1} \in \mathcal{K})$
– $M' \preceq_{K'} N' \;\Leftrightarrow M'^{\pi_0} \preceq_{K_0} N'^{\pi_0}, M'^{\pi_1} \preceq_{K_1} N'^{\pi_1}, (M'^{\delta} \preceq_K N'^{\delta})$

- *Remark:* $cat(\mathcal{K})_{strict}$ is the pullback in CAT, the category of all (large) categories.

- **Proposition:** $\mathcal{K}'$ is an AEC:

Proof. The validity of the axioms *order, isomorphism, coherence, reunion* follows from the relation $L' - str \cong (L_0 - str) \underset{L-str}{\times} (L_1 - str)$. For *Löwenhein-Skolem*: $LS(\mathcal{K}), LS(\mathcal{K}_0), LS(\mathcal{K}_1) \leqslant \lambda \Rightarrow LS(\mathcal{K}') \leqslant \lambda$. Take $X \subseteq M' \in \mathcal{K}'$ and consider the diagram:



$P' \in L' - str$ (well defined):
$(P')^0 \overset{h}{=} \bigcup_{n\in\mathbb{N}} P_{2n} \preceq_0 (M')^0$
$(P')^1 \cong \bigcup_{n\in\mathbb{N}} P_{2n+1} \preceq_1 (M')^1$
$(P')^1 \subseteq_{L_1} (M')^1$
iso condition in $\mathcal{K}_1$ entails $(P')^1 \preceq_1 (M')^1$
thus $P' \preceq_{K'} M'$, $X \subseteq P'$, $\sharp P' \leqslant \lambda$

- *Special case*:

$L = L_0 \cap L_1$, $L' = L_0 \cup L_1$ (pullback/pushout)



□

AECs for g-types

Similar to pullback (smaller class, but with pullback relation).

2.9. Given $M \in \mathcal{K}$, consider $\mathcal{K}_M$ the class of all pairs (N, f) where $N \in \mathcal{K}$, $f : M \rightarrowtail N \in cat(\mathcal{K})$.

- $\mathcal{K}_M \subseteq L(M) - str$
- binary relation: $(N_0, f_0) \preceq_{K_M} (N_1, f_1)$ iff $N_0 \preceq_K N_1$ and $(N_0, f_0) \subseteq_{L(M)} (N_1, f_1)$
- $(\mathcal{K}_M, \preceq_{K_M})$ is AEC $\quad (LS(\mathcal{K}) \leqslant \lambda \Rightarrow LS(\mathcal{K}_M) \leqslant \lambda + \sharp M)$
- $cat(\mathcal{K}_M)$: commutative triangles in $cat(\mathcal{K})$ over M



□

2.10. *An AEC for g-types*

Given $M \in \mathcal{K}$ and I a set, consider $\mathcal{K}_{M,I}$ the class of all triples $(N, f, \bar{a})$ where $N \in \mathcal{K}$, $\bar{a} : I \to N$, $f : M \rightarrowtail N \in cat(\mathcal{K})$.

- $\mathcal{K}_{M,I} \subseteq L(M \cup I) - str$
- binary relation: $(N_0, f_0, \bar{a}_0) \preceq_{K_{MI}} (N_1, f_1, \bar{a}_1)$ iff $N_0 \preceq_K N_1$ and $(N_0, f_0, \bar{a}_0) \subseteq_{L(M,I)} (N_1, f_1, \bar{a}_1)$
- $(\mathcal{K}_{M,I}, \preceq_{K_{M,I}})$ is AEC $\quad (LS(\mathcal{K}) \leqslant \lambda \Rightarrow LS(\mathcal{K}_{M,I}) \leqslant \lambda + \sharp M + \sharp I)$
- $cat(\mathcal{K}_{M,I})$: commutative triangles in $cat(\mathcal{K})$ over M and I
- we may assume $M \cap I = \emptyset$ (e.g., take $I := C \times \{M\}$)



□

Fact 2.11. *The I-types over M are precisely the g-maximal consistent theories in $\mathcal{K}_{M,I}$, thus they are AECs in the language $L(M \cup I)$.*

$$g - T(M, I) = Max(\mathcal{K}_{M,I})$$

□

Fact 2.12. Induced arrows on g-types:
$\alpha : L \to L'$, $\alpha^{\star}\restriction : \mathcal{K}' \to \mathcal{K}$, $h' : M_0' \to M_1' \in cat(\mathcal{K}')$, C a set of constants

- *a commutative diagram of languages and morphisms:*

– $I_j = C \times \{|M_j'|\}$, $j = 0, 1$;
– $M_j := M_j'^{\alpha} \Rightarrow |(M_j)^{\alpha}| = |M_j'|$, $j = 0, 1$;
– $h := (h')^{\alpha} : M_0 \to M_1$

$$\begin{array}{ccc} L(M_0 \cup I_0) & \xrightarrow{\widetilde{h}} & L(M_1 \cup I_1) \\ \downarrow{\scriptstyle \widetilde{\alpha}^0} & & \downarrow{\scriptstyle \widetilde{\alpha}^1} \\ L'(M'_0 \cup I_0) & \xrightarrow[\widetilde{h'}]{} & L'(M'_1 \cup I_1) \end{array}$$

- *a commutative diagram of AECs and morphisms:*

$$\begin{array}{ccc} \mathcal{K}_{M_0,I_0} & \xleftarrow{(\widetilde{h})^\star} & \mathcal{K}_{M_1,I_1} \\ \uparrow{\scriptstyle (\widetilde{\alpha}^0)^\star} & & \uparrow{\scriptstyle (\widetilde{\alpha}^1)^\star} \\ \mathcal{K}'_{M'_0,I_0} & \xleftarrow[(\widetilde{h'})^\star]{} & \mathcal{K}'_{M'_1,I_1} \end{array}$$

- *a commutative diagram of sets and functions:*

$$\begin{array}{ccc} Max(\mathcal{K}_{M_0,I_0}) & \xleftarrow{\bar{\widetilde{h}}} & Max(\mathcal{K}_{M_1,I_1}) \\ \uparrow{\scriptstyle \bar{\widetilde{\alpha}}^0} & & \uparrow{\scriptstyle \bar{\widetilde{\alpha}}^1} \\ Max(\mathcal{K}'_{M'_0,I_0}) & \xleftarrow[\bar{\widetilde{h'}}]{} & Max(\mathcal{K}'_{M'_1,I_1}) \end{array}$$

equivalently

$$\begin{array}{ccc} g-T(M_0,I_0) & \xleftarrow{\bar{\widetilde{h}}} & g-T(M_1,I_1) \\ \uparrow{\scriptstyle \bar{\widetilde{\alpha}}^0} & & \uparrow{\scriptstyle \bar{\widetilde{\alpha}}^1} \\ g-T(M'_0,I_0) & \xleftarrow[\bar{\widetilde{h'}}]{} & g-T(M'_1,I_1) \end{array}$$

□

Fact 2.13. λ-Topology on g-types (II)*:*

(i) $\lambda \geqslant LS(\mathcal{K}) \Rightarrow X^\lambda(M,I)$ *is the least topology such that:*
– $\widetilde{i_N} : g-T(M,I) \to g-T(N,I)$ *is continuous,* $N \in S^\lambda(M) := \{N \in \mathcal{K} : N \preceq_K M, \sharp N \leqslant \lambda\}$, $X^\lambda(N,I) = (g-T(N,I), discrete)$
equivalently:
– $can^\lambda_M : X^\lambda(M,I) \to lim_{N \in S^\lambda(M)} X^\lambda(N,I)$ *(with induced pro-topology) is continuous;*
equivalently:
– *if* $M \cong colim_{j \in J} P_j$ *(directed colimit),* $\sharp P_j \leqslant \lambda$
$can : X^\lambda(M,I) \to lim_{j \in J} X^\lambda(P_j,I) \approx lim_{N \in S^\lambda(M)} X^\lambda(N,I)$ *(with induced pro-topology) is continuous*

(ii) Properties of $\mathcal{K}$*: pro-topology ?, injective?, surjective?*
–λ*-Tame* $\leftrightsquigarrow \forall M,I$*, can is injective* (T_2)
–"λ*-compact*" $\leftrightsquigarrow \forall M,I$*, can is surjective*

(iii) Induced arrows on g-types:
$\alpha : L \to L'$, $\alpha^\star \restriction : \mathcal{K}' \to \mathcal{K}$, $h' : M'_0 \to M'_1 \in cat(\mathcal{K}')$, C *a set of constants,* $\lambda \geqslant LS(\mathcal{K}), LS(\mathcal{K}')$

- $M' \cong colim_{j \in J} P'_j \Rightarrow M'^\alpha \cong colim_{j \in J} P'^\alpha_j$

$\hat{h}' : S^\lambda(M'_0) \to S^\lambda(M'_1)$

- *two diagrams of spaces and continuous functions*



then



- *If $\mathcal{K}'$ has* **(AP)** $\Rightarrow \bar{\tilde{h}}' : X^\lambda(M_1', I) \twoheadrightarrow X^\lambda(M_0', I)$ *and* $\bar{\tilde{h}} : X^\lambda(M_1, I) \twoheadrightarrow X^\lambda(M_0, I)$ *are surjective.* □

2.2 New (global) axioms and applications

"Local" axioms and applications:

Where "local" means "for each AEC fixed"
Ex.:
(JEP): Joint Embbeding Property;
(AP): Amalgamation Property;
(LRP): "Local Robinson's Property": $M_0 \equiv_{\mathcal{K}} M_1$ iff $\exists N \in \mathcal{K} \exists j_0, j_1 \in cat(\mathcal{K})\ j_0 : M_0 \rightarrowtail N$, $j_1 : M_1 \rightarrowtail N$

Robinson's diagram method:

2.14. If $\mathcal{K}$ is an AEC in language L, then for each set of "new constants" $L' := L \cup \{c_i : i \in I\}$ it is available the AEC $\mathcal{K}'$ in the language L' whose objects are pairs $(M, \bar{a})$, where $\bar{a}$ is a I-sequence of elements of M and $(M, \bar{a}) \preceq_{\mathcal{K}'} (N, \bar{b})$ iff: $M \preceq_{\mathcal{K}} N$ and $\bar{a} = \bar{b}$ $(\Leftrightarrow\ M \preceq_{\mathcal{K}} N$ and $(M, \bar{a}) \subseteq_{L'} (N, \bar{b}))$

- $\mathcal{K}'$ it is the pullback:

□

Proposition 2.15.(LRP) $\exists j : M \rightarrowtail N \in cat(\mathcal{K})$ *iff* $\exists \bar{b}$ *such that* $(M, \bar{a}) \equiv_{K(M)} (N, \bar{b})$ *where* $\mathcal{K}(M) \subseteq L(M) - str$



$$(M, \bar{a}) \overset{t\restriction}{\underset{\cong}{\rightarrow}} (M', \bar{c}) \overset{incl}{\underset{\preceq}{\rightarrow}} (P, \bar{c})$$

$$(N, \bar{b}) \overset{h\restriction}{\underset{\cong}{\rightarrow}} (N', \bar{c}) \overset{incl}{\underset{\preceq}{\rightarrow}} (P, \bar{c})$$



$$M, \bar{a}) \overset{t\restriction}{\underset{\cong}{\to}} (M', \bar{c}) \overset{incl}{\underset{\prec}{\to}} (N', \bar{c}) \overset{(h\restriction)^{-1}}{\underset{\cong}{\to}} (N, \bar{b})$$

$$(M, \bar{a}) \overset{j}{\rightarrowtail} (N, \bar{b})$$

□

The (global) category AEC:

The Grothendieck's gluying of all AECs

objects: pairs $(M, \mathcal{K})$, where $M \in \mathcal{K}$

arrows: pairs $(h, \varphi) : (M, \mathcal{K}) \to (M', \mathcal{K}')$
$\varphi : L \to L'$, $\Phi = \varphi^\star \restriction : \mathcal{K}' \to \mathcal{K}$, $h : M \rightarrowtail \Phi(M') \in cat(\mathcal{K})$

identities: $(id_M, id_L) : (M, \mathcal{K}) \to (M, \mathcal{K})$

composition: $(M, \mathcal{K}) \overset{(h,\varphi)}{\to} (M', \mathcal{K}') \overset{(h',\varphi')}{\to} (M'', \mathcal{K}'') = (M, \mathcal{K}) \overset{(\Phi(h')\circ h, \varphi'\circ\varphi)}{\to} (M'', \mathcal{K}'')$

$(M \overset{h}{\rightarrowtail} \Phi(M') \overset{\Phi(h')}{\rightarrowtail} \Phi(\Phi'(M'')))$ □

2.16. *New (global) equivalences in AEC*:

– the connection relation: $(N_0, \mathcal{K}_0) \equiv (N_1, \mathcal{K}_1)$

– relative equivalence over $M \in \mathcal{K}$:
$(N_0, \mathcal{K}_0) \equiv_{(M,\mathcal{K})} (N_1, \mathcal{K}_1)$, with $h_i : M \rightarrowtail (N_i)^{\alpha_i}$

Ex. I-types over M : $(M, \mathcal{K}) \to ((N, \bar{a}), \mathcal{K}(I))$ □

New (global) axioms in AEC

Goal: Determine interesting subcategories of AEC satisfying interesting (global) axioms.

(GAP) "Global Amalgamation Property"
entails a simpler description of the relative equivalence over $M \in \mathcal{K}$:
$(N_0, \mathcal{K}_0) \equiv_{(M,\mathcal{K})} (N_1, \mathcal{K}_1)$, with $h_i : M \rightarrowtail (N_i)^{\alpha_i}$
iff $\exists \mathcal{K}' \exists P' \exists j_i : N_i \rightarrowtail P'^{\beta_i}$ such that the diagram below commutes



(TRP) "Transversal Robinson Property":
• $\alpha_i : L \to L_i$, $i = 0, 1$
$\mathcal{K}, \mathcal{K}_0, \mathcal{K}_1$

• $(M_0)^K \equiv_K (M_1)^K \Rightarrow \exists N_0 \in \mathcal{K}_0$ such that:
– $M_0 \prec_{K_0} N_0$ in $\mathcal{K}_0$
– $\exists j : (M_1)^K \rightarrowtail (N_0)^K$ in $cat(\mathcal{K})$

(GRP) "Global Robinson Property":
• $\alpha_i : L \to L_i$, $\beta_i : L_i \to (L_0 \cup L_1)/\sim$, $i = 0, 1$
$\mathcal{K}, \mathcal{K}_0, \mathcal{K}_1$, $\mathcal{K}' = pullback$

• $(M_0)^K \equiv_K (M_1)^K \Rightarrow \exists P' \in \mathcal{K}'$ such that:
– $M_0 \prec_{K_0} (P')^{K_0}$ in $\mathcal{K}_0$
– $\exists h : M_1 \rightarrowtail (P')^{K_1}$ in $cat(\mathcal{K}_1)$

Proposition 2.17. *(TRP) ⇒ (GRP)*
Proof. *Just take zig-zags as in the proof of LS property in the pullback AEC* □



$P' \in L' - str$ (well defined):
$(P')^0 = \bigcup_{n\in\mathbb{N}} P_{2n} \in \mathcal{K}_0$
$(P')^1 \overset{h}{\cong} \bigcup_{n\in\mathbb{N}} P_{2n+1} \in \mathcal{K}_1$
thus $P' \in \mathcal{K}'$ and
– $M_0 \preceq_{K_0} (P')^{K_0}$ in $\mathcal{K}_0$
– $\exists h : M_1 \rightarrowtail (P')^{K_1}$ in $cat(\mathcal{K}_1)$

Craig Interpolation Property:

2.18. *Craig interpolation in FOL*:

- $\psi_0, \psi_1 \in L' - sent$ $\psi_0 \vdash \psi_1 \Rightarrow$ exists $\theta \in L = L_0 \cap L_1$: $\psi_0 \vdash \theta$, $\theta \vdash \psi_1$
- $\{\psi_0, \neg\psi_1\} \vdash \bot \Rightarrow$ exists $\theta \in L = L_0 \cap L_1$: $\psi_0 \vdash \theta$, $\neg\psi_1 \vdash \neg\theta$
- $T_0 \cup T_1$ has no L'-model $\Rightarrow$ exists $\theta \in L = L_0 \cap L_1$: $T_0 \vDash \theta$, $T_1 \vDash \neg\theta$
- $L = L_0 \cap L_1, T_i \restriction_L = \{\psi \in L - sent : T_i \vdash_L \psi\}$

$T_0 \cup T_1$ has no L'-model $\Rightarrow$ $T_0 \restriction_L \cup T_1 \restriction_L$ has no L-model

- $\nexists M' \in L' - str$ such that $(M')^{L_i} \in Mod(T_i)$, $i = 0, 1$ $\Rightarrow$ $\nexists M \in L - str$ with $M \in Mod(T_0\restriction_L) \cap Mod(T_1\restriction_L)$ □

(CIP) "Craig Interpolation Property":

- $\alpha_i : L \to L_i$, $\beta_i : L_i \to (L_0 \cup L_1)/\sim$, $i = 0, 1$

$\mathcal{K}, \mathcal{K}_0, \mathcal{K}_1, \mathcal{K}'$ = *pullback*
$Mod(T_i) \subseteq \mathcal{K}_i$, $i = 0, 1$.

- $\nexists M' \in \mathcal{K}'$ such that $(M')^{K_i} \in Mod(T_i)$, $i = 0, 1$ $\Rightarrow$ $\nexists M \in \mathcal{K}$ with $M \in Mod(T_0\restriction_K) \cap Mod(T_1\restriction_K)$ (the induced theories)

Proposition 2.19. $(TRP) \Rightarrow (CIP)$
Proof. *(contrapositive)*

Let $N \in \mathcal{K}$ with $N \in Mod(T_0{\restriction}_L) \cap Mod(T_1{\restriction}_L)$.
Then: there are $M_i \in Mod(T_i)$, $i = 0, 1$, with $(M_0)^K \equiv_K N \equiv_K (M_1)^K$.
As $(TRP) \Rightarrow (GRP)$, there is $P' \in \mathcal{K}'$ such that:
- $M_0 \preceq_{K_0} (P')^{K_0}$ *in* $\mathcal{K}_0$
- $\exists h : M_1 \rightarrowtail (P')^{K_1}$ *in* $cat(\mathcal{K}_1)$.

Then:
- $(P')^{K_0} \equiv_{K_0} M_0 \in Mod(T_0)$;
- $(P')^{K_1} \equiv_{K_1} M_1 \in Mod(T_1)$. □

3 Final Remarks and Future Works

- Introduce and study other global axioms (inspirated on model theory of FOL).
- Consider functors induced by more general language morphisms $f : L \to L'$:

(i) non injective;
(ii) "flexible morphism":
$f_n \in L$ (n-nary functional symbol) $\mapsto t_n \in L'$ (n-ary term)
$R_n \in L$ (n-ary relational symbol) $\mapsto \varphi_n \in L'$ (n-ary atomic formula)

- Consider more general functors between AECs (ex. functors that are not "over Set").
- Study metric abstract elementary classes (MAECs) in this global setting.

References

[**AR**] J. Adámek, J. Rosický, **Locally Presentable and Accessible Categories**, Lecture Notes Series of the LMS **189**, Cambridge University Press, Cambridge, Great Britain, 1994.

[**Bal1**] J. Baldwin, *Abstract Elementary Classes: Some Answers, More Questions*, July 7 2006, 18 pages.

[**Bal2**] J. Baldwin, **Categoricity**, University Lecture Series **50**, American Mathematical Society, 2009.

[**BR**] T. Beke, J. Rosický, *Abstract elementary classes and accessible categories*, Annals of Pure and Applied Logic **163 (12)**, 2012, 2008-2017.

[**Kue**] D. Kueker, *Abstract elementary classes and infinitary logics* , Annals of Pure and Applied Logic, **156(11)**, 2008, 274-286.

[**Lie1**] M.J. Lieberman, **Topological and Category-Theoretic Aspects of Abstract Elementary Classes**, Phd, University of Michigan, 99 pages, 2009.

[**Lie2**] M.J. Lieberman, *Category-theoretic aspects of of abstract elementary classes*, Annals of Pure and Applied Logic, **162(11)**, 2011, 903-915.

[**She**] S. Shelah, *Classification of nonelementary classes II, abstract elementary classes*, Lecture Notes in Mathematics **1292**, 1987, 419-497.

[**Van**] M. VanDieren, *Categoricity in Abstract Elementary Classes with no maximal models*, availabel at *h*ttp://arxiv.org/pdf/math/0510579.pdf, 2005, 57 pages.